## **ERGON**



## IMPROVES RESISTANCE TO STRESS IMPROVES PHOTOSYNTHETIC EFFICIENCY ALLOWED IN ORGANIC FARMING

ERGON is a solution that combines:

- the biostimulating activity of Ascophyllum nodosum seaweed extract, thanks to its content of vitamins, polysaccharides and natural growth promoters
- the energizing activity of the free amino acids derived from epithelium's enzymatic hydrolysis
- the greening activity of iron that being a key element in chlorophyll bio-synthesis, improves also the photosynthetic process ERGON, readily absorbed thanks to its acidic pH, keeps all the biochemical pathways active. It is therefore recommended on all crops going through either delays or interruptions in vegetative growth caused by stress conditions and whenever the crop is under stress due to high production targets.

| CROP  | TIME OF APPLICATION  | DOSE/HECTARE* |
|---|--|---------------|
| Kiwifruit e Grapes  | 2-3 applications from vegetative restart to fruit enlargement, every 10-15 days        | 2-3 kg        |
| Citrus (Orange, Bergamot, Clementine, Lemon, Tangerine)   | 2-3 applications from vegetative restart to fruit enlargement, every 10-15 days        | 2-3 kg        |
| Stone fruits (Apricot, Cherry, Nectarine, Peach, Plum) e Pome fruits (Quince, Apple, Pear)                | 2-3 applications from vegetative restart to fruit enlargement, every 10-15 days        | 2-3 kg        |
| Strawberries  | 3 applications from the beginning of flowering, every 10-15 days                       | 2-3 kg        |
| Fruiting vegetables (Watermelon, Cucumber, Eggplant, Melon, Pepper, Tomato, Zucchini, Pumpkin)            | 3-4 applications from the beginning of flowering, every 10-15 days                     | 2-3 kg        |
| Flowers and ornamentals   | 3-4 applications from the beginning of flowering, every 10-15 days                     | 2-3 kg        |
| Leafy vegetables (Chicory, Lettuce, Radicchio, Rocket, Escarole, Celery, Spinach)                         | carole, At early vegetative phases or whenever a prompt vegetative restart is required |               |
| Cereal crops (Oats, Wheat, Corn, Barley, Rice, Rye, Sorghum, Triticale)                                   | At early vegetative phases in combination with herbicide/pesticide treatments          | 3 kg          |
| Industrial crops (Beets, Sugarcane, Rapeseed, Cotton,<br>Sunflower, Industrial tomato, Soybeans, Tobacco) | At early vegetative phases or whenever a prompt vegetative restart is required         | 3 kg          |

| COMPOSITION                     |        |  |
|---------------------------------|--------|--|
| Organic nitrogen (N)            | 4.00%  |  |
| Organic nitrogen (N) soluble    | 3.60%  |  |
| Carbon (C) of biological origin | 12.00% |  |
| Iron (Fe) soluble in water      | 3.00%  |  |

| PHYSICO-CHEMICAL FEATURES       |                      |  |
|---------------------------------|----------------------|--|
| LIQUID                          |                      |  |
| pH (sol 1%)                     | 2.90                 |  |
| Conductivity E.C. S/cm (1‰)     | 505                  |  |
| Density (g/cm³)/Specific weight | 1.10                 |  |
| METHOD OF USE                   | <b>₽</b>             |  |
|                                 | Foliar fertilization |  |

**PACKAGING: 1 - 6 - 25 KG** 

**NOTE**: ERGON can be also used in fertigation, with a 1,5-3 kg/1000 m<sup>2</sup> dose and every 10-12 days.